

PROBABILITY AND STATISTICS




SUBRATA SAHA

AGENDA

—

INDEPENDENT EVENTS



CONDITIONAL PROBABILITY

3 INDEPENDENT EVENTS

Giving more information can change the probability of an event.

EXAMPLE :

If a coin is tossed two times then what is the probability of two Heads?

ANSWER : $\frac{1}{4}$.

EXAMPLE :

If a coin is tossed two times then what is the probability of two Heads, *given that the first toss gave Heads* ?

ANSWER : $\frac{1}{2}$.

NOTE :

Several examples will be about *playing cards* .

A standard *deck* of *playing cards* consists of 52 cards :

- Four *suits* :

Hearts , Diamonds (*red*) , and Spades , Clubs (*black*) .

- Each suit has 13 cards, whose *denomination* is

2 , 3 , \dots , 10 , Jack , Queen , King , Ace .

- The Jack , Queen , and King are called *face cards* .

EXERCISE :

Suppose we draw a card from a shuffled set of 52 playing cards.

- What is the probability of drawing a Queen ?
- What is the probability of drawing a Queen, given that the card drawn is of *suit* Hearts ?
- What is the probability of drawing a Queen, given that the card drawn is a *Face card* ?

What do the answers tell us?

(We'll soon learn the events "Queen" and "Hearts" are *independent*.)

CONDITIONAL PROBABILITY

6 INDEPENDENT

EVENTS

The two preceding questions are examples of *conditional probability*

Conditional probability is an *important* and *useful* concept.

If E and F are events, *i.e.*, subsets of a sample space \mathcal{S} , then

$P(E|F)$ *is the conditional probability of E , given F ,*

defined as

$$P(E|F) \equiv \frac{P(EF)}{P(F)} .$$

or, equivalently

$$P(EF) = P(E|F) P(F) ,$$

(assuming that $P(F)$ is not zero).



CONDITIONAL PROBABILITY

7 INDEPENDENT

EVENTS

28

$$P(E|F) \equiv \frac{P(EF)}{P(F)}$$



Suppose that the 6 outcomes in \mathcal{S} are equally likely.

What is $P(E|F)$ in each of these two cases ?



CONDITIONAL PROBABILITY

8 INDEPENDENT

EVENTS

28

$$P(E|F) \equiv \frac{P(EF)}{P(F)}$$



Suppose that the 6 outcomes in \mathcal{S} are equally likely.

What is $P(E|F)$ in each of these two cases ?

EXAMPLE : Suppose a coin is tossed two times.

The sample space is

$$\mathcal{S} = \{HH, HT, TH, TT\}.$$

Let E be the event "*two Heads*", i.e.,

$$E = \{HH\}.$$

Let F be the event "*the first toss gives Heads*", i.e.,

$$F = \{HH, HT\}.$$

Then

$$EF = \{HH\} = E \quad (\text{since } E \subset F).$$

We have

$$P(E|F) = \frac{P(EF)}{P(F)} = \frac{P(E)}{P(F)} = \frac{\frac{1}{4}}{\frac{2}{4}} = \frac{1}{2}.$$

EXAMPLE :

Suppose we draw a card from a shuffled set of 52 playing cards.

- What is the probability of drawing a Queen, given that the card drawn is of *suit* Hearts ?

ANSWER :

$$P(Q|H) = \frac{P(QH)}{P(H)} = \frac{\frac{1}{52}}{\frac{13}{52}} = \frac{1}{13} .$$

- What is the probability of drawing a Queen, given that the card drawn is a *Face card* ?

ANSWER :

$$P(Q|F) = \frac{P(QF)}{P(F)} = \frac{P(Q)}{P(F)} = \frac{\frac{4}{52}}{\frac{12}{52}} = \frac{1}{3} .$$

(Here $Q \subset F$, so that $QF = Q$.)

The probability of an event E is sometimes computed more easily
if we condition E on another event F ,

namely, from

$$\begin{aligned} P(E) &= P(E(F \cup F^c)) \quad (\text{Why ?}) \\ &= P(EF \cup EF^c) = P(EF) + P(EF^c) \quad (\text{Why ?}) \end{aligned}$$

and

$$P(EF) = P(E|F) P(F) \quad , \quad P(EF^c) = P(E|F^c) P(F^c) \quad ,$$

we obtain this *basic formula*

$$P(E) = P(E|F) \cdot P(F) + P(E|F^c) \cdot P(F^c) .$$

EXAMPLE :

An insurance company has these data :

The probability of an insurance claim in a period of one year is

4 percent for persons under age 30

2 percent for persons over age 30

and it is known that

30 percent of the targeted population is under age 30.

What is the probability of an insurance claim in a period of one year for a randomly chosen person from the targeted population?

SOLUTION :

Let the sample space \mathcal{S} be all persons under consideration.

Let C be the event (subset of \mathcal{S}) of persons filing a claim.

Let U be the event (subset of \mathcal{S}) of persons under age 30.

Then U^c is the event (subset of \mathcal{S}) of persons over age 30.

Thus

$$\begin{aligned} P(C) &= P(C|U) P(U) + P(C|U^c) P(U^c) \\ &= \frac{4}{100} \frac{3}{10} + \frac{2}{100} \frac{7}{10} \\ &= \frac{26}{1000} = 2.6\% . \end{aligned}$$

EXAMPLE :

Two balls are drawn from a bag with 2 *white* and 3 *black* balls.

There are 20 outcomes (*sequences*) in \mathcal{S} . (**Why ?**)

What is the probability that *the second ball is white* ?

SOLUTION :

Let F be the event that *the first ball is white*.

Let S be the event that *the second second ball is white*.

Then

$$P(S) = P(S|F) P(F) + P(S|F^c) P(F^c) = \frac{1}{4} \cdot \frac{2}{5} + \frac{2}{4} \cdot \frac{3}{5} = \frac{2}{5}.$$

QUESTION : Is it surprising that $P(S) = P(F)$?

EXAMPLE : (continued ...)

Is it surprising that $P(S) = P(F)$?

ANSWER : Not really, if one considers the sample space \mathcal{S} :

$$\left\{ \begin{array}{llll} \mathbf{w}_1 \mathbf{w}_2, & \mathbf{w}_1 b_1, & \mathbf{w}_1 b_2, & \mathbf{w}_1 b_3, \\ \mathbf{w}_2 \mathbf{w}_1, & \mathbf{w}_2 b_1, & \mathbf{w}_2 b_2, & \mathbf{w}_2 b_3, \\ b_1 \mathbf{w}_1, & b_1 \mathbf{w}_2, & b_1 b_2, & b_1 b_3, \\ b_2 \mathbf{w}_1, & b_2 \mathbf{w}_2, & b_2 b_1, & b_2 b_3, \\ b_3 \mathbf{w}_1, & b_3 \mathbf{w}_2, & b_3 b_1, & b_3 b_2 \end{array} \right\},$$

where outcomes (*sequences*) are assumed equally likely.

EXAMPLE :

Suppose we draw *two cards* from a shuffled set of 52 playing cards.

What is the probability that the second card is a Queen ?

ANSWER :

$$P(2^{\text{nd}} \text{ card } Q) =$$

$$P(2^{\text{nd}} \text{ card } Q | 1^{\text{st}} \text{ card } Q) \cdot P(1^{\text{st}} \text{ card } Q)$$

$$+ P(2^{\text{nd}} \text{ card } Q | 1^{\text{st}} \text{ card not } Q) \cdot P(1^{\text{st}} \text{ card not } Q)$$

$$= \frac{3}{51} \cdot \frac{4}{52} + \frac{4}{51} \cdot \frac{48}{52} = \frac{204}{51 \cdot 52} = \frac{4}{52} = \frac{1}{13}.$$

QUESTION : Is it surprising that $P(2^{\text{nd}} \text{ card } Q) = P(1^{\text{st}} \text{ card } Q)$?

A useful formula that "*inverts conditioning*" is derived as follows :

Since we have both

$$P(EF) = P(E|F) P(F) ,$$

and

$$P(EF) = P(F|E) P(E) .$$

If $P(E) \neq 0$ then it follows that

$$P(F|E) = \frac{P(EF)}{P(E)} = \frac{P(E|F) \cdot P(F)}{P(E)} ,$$

and, using the earlier useful formula, we get

$$P(F|E) = \frac{P(E|F) \cdot P(F)}{P(E|F) \cdot P(F) + P(E|F^c) \cdot P(F^c)} ,$$

which is known as *Bayes' formula* .

EXAMPLE : Suppose 1 in 1000 persons has a certain disease.

A test detects the disease in 99 % of diseased persons.

The test also "detects" the disease in 5 % of healthy persons.

With what probability does a positive test diagnose the disease?

SOLUTION : Let

$D \sim$ "diseased" , $H \sim$ "healthy" , $+$ \sim "positive".

We are given that

$$P(D) = 0.001 , \quad P(+|D) = 0.99 , \quad P(+|H) = 0.05 .$$

By Bayes' formula

$$\begin{aligned} P(D|+) &= \frac{P(+|D) \cdot P(D)}{P(+|D) \cdot P(D) + P(+|H) \cdot P(H)} \\ &= \frac{0.99 \cdot 0.001}{0.99 \cdot 0.001 + 0.05 \cdot 0.999} \cong 0.0194 \quad (!) \end{aligned}$$

EXERCISE :

Suppose 1 in 100 products has a certain defect.

A test detects the defect in 95 % of defective products.

The test also "detects" the defect in 10 % of non-defective products.

- With what probability does a positive test diagnose a defect?

EXERCISE :

Suppose 1 in 2000 persons has a certain disease.

A test detects the disease in 90 % of diseased persons.

The test also "detects" the disease in 5 % of healthy persons.

- With what probability does a positive test diagnose the disease?

More generally, if the sample space \mathcal{S} is *the union of disjoint events*

$$\mathcal{S} = F_1 \cup F_2 \cup \dots \cup F_n ,$$

then for any event E

$$P(F_i|E) = \frac{P(E|F_i) \cdot P(F_i)}{P(E|F_1) \cdot P(F_1) + P(E|F_2) \cdot P(F_2) + \dots + P(E|F_n) \cdot P(F_n)}$$

EXERCISE :

Machines M_1, M_2, M_3 produce these *proportions* of a article

Production : $M_1 : 10 \% , \quad M_2 : 30 \% , \quad M_3 : 60 \% .$

The probability the machines produce *defective* articles is

Defects : $M_1 : 4 \% , \quad M_2 : 3 \% , \quad M_3 : 2 \% .$

What is the probability a random article was made by machine M_1 , given that it is defective?

Independent Events

Two events E and F are *independent* if

$$P(EF) = P(E) P(F) .$$

In this case

$$P(E|F) = \frac{P(EF)}{P(F)} = \frac{P(E) P(F)}{P(F)} = P(E) ,$$

(assuming $P(F)$ is not zero).

Thus

knowing F occurred doesn't change the probability of E .

EXAMPLE : Draw *one* card from a deck of 52 playing cards.

Counting outcomes we find

$$P(\text{Face Card}) = \frac{12}{52} = \frac{3}{13} ,$$

$$P(\text{Hearts}) = \frac{13}{52} = \frac{1}{4} ,$$

$$P(\text{Face Card and Hearts}) = \frac{3}{52} ,$$

$$P(\text{Face Card}|\text{Hearts}) = \frac{3}{13} .$$

We see that

$$P(\text{Face Card and Hearts}) = P(\text{Face Card}) \cdot P(\text{Hearts}) \quad (= \frac{3}{52}) .$$

Thus the events "*Face Card*" and "*Hearts*" are *independent*.

Therefore we also have

$$P(\text{Face Card}|\text{Hearts}) = P(\text{Face Card}) \quad (= \frac{3}{13}) .$$

CONDITIONAL PROBABILITY

22 INDEPENDENT

EVENTS

28

EXERCISE :

CONDITIONAL PROBABILITY

23 INDEPENDENT EVENTS

Which of the following pairs of events are independent?

- (1) drawing "Hearts" and drawing "Black" ,
- (2) drawing "Black" and drawing "Ace" ,
- (3) the event $\{2, 3, \dots, 9\}$ and drawing "Red" .

EXERCISE : *Two* numbers are drawn at random from the set
 $\{ 1 , 2 , 3 , 4 \} .$

If *order is not important* then what is the sample space \mathcal{S} ?

Define the following functions on \mathcal{S} :

$$X(\{i, j\}) = i + j , \quad Y(\{i, j\}) = |i - j| .$$

Which of the following pairs of events are independent?

$$(1) \quad X = 5 \quad \text{and} \quad Y = 2 ,$$

$$(2) \quad X = 5 \quad \text{and} \quad Y = 1 .$$

REMARK :

X and Y are examples of *random variables* . (More soon!)

EXAMPLE : If E and F are *independent* then so are E and F^c .

PROOF : $E = E(F \cup F^c) = EF \cup EF^c$, where

EF and EF^c are *disjoint* .

Thus

$$P(E) = P(EF) + P(EF^c) ,$$

from which

$$P(EF^c) = P(E) - P(EF)$$

$$= P(E) - P(E) \cdot P(F) \quad (\text{since } E \text{ and } F \text{ independent})$$


$$= P(E) \cdot (1 - P(F))$$

$$= P(E) \cdot P(F^c) .$$


EXERCISE :

Prove that if E and F are *independent* then so are E^c and F^c .

NOTE : *Independence* and *disjointness* are different things !



Independent, but not disjoint.



Disjoint, but not independent.

(The six outcomes in S are assumed to have equal probability.)

If E and F are *independent* then $P(EF) = P(E) P(F)$.

If E and F are *disjoint* then $P(EF) = P(\emptyset) = 0$.

If E and F are *independent and disjoint* then one has *zero probability* !

CONDITIONAL PROBABILITY

26 INDEPENDENT
EVENTS

28

Three events E , F , and G are *independent* if

$$P(EFG) = P(E) P(F) P(G) .$$

and

$$P(EF) = P(E) P(F) .$$

$$P(EG) = P(E) P(G) .$$

$$P(FG) = P(F) P(G) .$$

EXERCISE : Are the three events of drawing

- (1) a red card ,
- (2) a face card ,
- (3) a Heart or Spade ,

independent ?

EXERCISE :

A machine M consists of three *independent parts*, M_1 , M_2 , and M_3 .

Suppose that

M_1 functions properly with probability $\frac{9}{10}$,

M_2 functions properly with probability $\frac{9}{10}$,

M_3 functions properly with probability $\frac{8}{10}$,

and that

the machine M functions if and only if *its three parts function*.

- What is the probability for the machine M to *function* ?
- What is the probability for the machine M to *malfunction* ?



THANK YOU

SUBRATA SAHA
SUBRATAISTATAMIKARANA.CO.IN